

Conclusions

Warfare has long been recognised as a stimulus for advancing surgical and medical care. However treatment of maxillofacial injuries sustained in Afghanistan has required the use of both traditional and contemporary techniques to achieve the best possible outcome for wounded British servicemen evacuated to RCDM. As the experience of the maxillofacial team in managing these injuries develops, further refinements will undoubtedly follow but it is likely that there will always be a role for these traditional techniques.

Acknowledgments

The authors would like to thank the following for their assistance in the management of these complex patients and the preparation of this manuscript: Professor M Midwinter, Defence Professor of Surgery and head of the Academic Department of Military Surgery and Trauma, Mr I Sharp, Mr R Williams, Mr S Dover, Mr K Webster, Mr B Speculand.

References

1. Breeze J, Gibbons AJ, Opie NJ, Monaghan A. Maxillofacial injuries in military personnel treated at the Royal Centre for Defence Medicine June 2001 to December 2007. *Br J Oral Maxillofac Surg*. 2009 Nov 6. (Epub ahead of print)
2. Dobson JE, Newell MJ, Shepherd JP. Trends in Maxillofacial Injuries in Wartime (1914-1986). *Br J Oral Maxillofac Surg* 1989; 27: 441-450.
3. Lew TA, Walker JA, Wenke JC, Blackburne LH, Hale RG. Characterization of Craniomaxillofacial Battle Injuries Sustained by United States Service Members in the Current Conflicts of Iraq and Afghanistan. *J Oral Maxillofac Surg* 2010;68:3-7
4. Scott PT, Peterson K, Fishbain J et al. Acinetobacter baumannii infections amongst patients at military medical facilities treating injured US service members, 2002- 2004. *MMWR Morb Mortal Wkly Rep* 2004; 53 (45): 1063- 1066.
5. Tien HC, Battad A, Bryce EA et al. Multi-drug resistant Acinetobacter infections in critically injured Canadian forces soldiers. *BMC Infectious Diseases* 2007; 7: 95.
6. Powers DB, Will MJ, Bourgeois SL, Hatt HD. Maxillofacial Trauma Treatment Protocol. *Oral Maxillofac Surg Clin N Am* 2005; 17: 341- 355.
7. Futran ND. Maxillofacial Trauma Reconstruction. *Facial Plast Surg Clin N Am*; 17 (2009):239-251.
8. Futran ND, Farwell DG, Smith RB et al. Definitive management of severe facial trauma utilizing free tissue transfer. *Otolaryngol Head and Neck Surg* 2005;132:75-85.
9. Vishal S, Farwell DG. Gunshot wounds to the head and neck. *Curr Opin Otolaryngol Head and Neck Surg* 2007;15:213 -218.
10. Tan YH, Zhou SX, Liu YQ, Liu BL, Li ZY. Small-vessel pathology and anastomosis following maxillofacial firearm wounds: an experimental study. *J Oral Maxillofac Surg* 1991; 49(4): 348-52
11. Ellis E 3rd, Muniz O, Anand K. Treatment considerations for comminuted mandibular fractures. *J Oral Maxillofac Surg* 2003; 61(8): 861-70.

Lessons Learned in Oral and Maxillofacial Surgery from British Military Deployments in Afghanistan

Gibbons AJ¹, Mackenzie N²

¹Department of Oral and Maxillofacial Surgery, Ministry of Defence Hospital Unit, Peterborough Hospital, Thorpe Road, Peterborough; ²Department of Oral and Maxillofacial Surgery, Ministry of Defence Hospital Unit, Queen Alexandra Hospital, Portsmouth, Hampshire, PO6 3LY

Abstract

This article describes the combined lessons learned from two deployments of a cadre of British Oral and Maxillofacial surgeons to Kandahar between July 2006 to April 2007, and September 2008 to April 2009.

Introduction

The British Oral and Maxillofacial Surgery Cadre have been deployed twice to the Role 3 multinational military hospital in Kandahar Afghanistan. Eight consultants deployed from July 2006 to April 2007, and five consultants, two of whom were accompanied by registrars, from September 2008 to April 2009. The British Military Oral and Maxillofacial Cadre holds 2 meetings a year at which cases are presented and experiences are shared. This article aims to outline the key surgical principles of oral and maxillofacial surgery as practiced in a conflict zone and to highlight some of the lessons learned by surgeons.

Work in the Role 3 Hospital

In the Role 3 hospital at Kandahar, all hospital staff are available on-call 24 hours a day. Clinical governance is implemented as it is in most developed nations' hospital systems. However, there is no need to meet targets and the ability to deliver patient centred care

without Western hospital management restrictions is, for many clinicians, a rewarding experience. Nevertheless, difficult patient management decisions are required. Like all hospitals, there is finite bed and operating theatre resources meaning there is an emphasis on moving patients on through the chain of evacuation so that these resources can be kept available. Mass casualty situations occur regularly and place demands on both beds and theatre time. Elective operating is done early in the morning to free theatres for later in the day when new casualties arrive. Military patients are frequently evacuated within 24 hours of injury following resuscitation and stabilization in the hospital. Afghan patients are passed back into the limited national military and civilian hospital network when deemed appropriate. Approximately one third of patients are NATO or coalition personnel and contracted civilians, one third Afghan army and police and one third Afghan civilians.

Military maxillofacial surgeons may be required to have extended competencies in ophthalmic surgery and neurosurgery. Evisceration, enucleation or partial exenteration of eyes for penetrating trauma is frequently required. In addition, a good knowledge of paediatric maxillofacial trauma is important. Teamwork with colleagues is essential and joint operating, assisting or being assisted by other surgeons, is often necessary.

Corresponding Author: Wing Commander A Gibbons, Consultant, Department of Oral and Maxillofacial Surgery, Ministry of Defence Hospital Unit, Peterborough Hospital, Thorpe Road, Peterborough, Cambridgeshire PE3 6DA
Email: Andrew.gibbons@pbh-tr.nhs.uk

Injury Patterns

A conflict zone hospital is a unique working environment. The challenges faced in treating military and civilian patients, including children, in a relatively austere environment should not be underestimated. Although surgeons can prepare by working in busy Western trauma centres, they will rarely treat blast or high energy transfer velocity ballistic injuries. The majority of ballistic injuries seen in the West are from low muzzle velocity hand guns. In recent military conflicts 75% of injuries are due to explosive mechanisms [1]. In the second Oral and Maxillofacial Surgery Cadre deployment in Afghanistan, over 50% of military oral and maxillofacial injuries were due to improvised explosive devices (IEDs) and the majority of gunshot wounds were from high energy rounds. Oral and Maxillofacial surgery patients with blast or other high energy transfer velocity injuries often present with extensive, contaminated soft tissue wounds and comminuted fractures. Tissue avulsion is frequently seen. Blast injuries may be further complicated by burns and multiple injuries both to the face and to other peripheral body sites. The use of body armour has led to a proportional increase in head and neck injuries [2].

Assessment

The Oral and Maxillofacial injury assessment should be a systematic and meticulous clinical examination between the top of the scalp and the clavicles. Frequently missed injuries are scalp lacerations, nasal fractures and small but deep penetrating shrapnel injuries. The penetrating and avulsive type of trauma seen is different to the blunt trauma seen in the West. A practical classification of wounds is to divide them into four groups (Box 1).

1. Soft tissue damage such as contusions, abrasions and lacerations.
2. Hard tissue damage i.e. fractures.
3. Soft tissue loss.
4. Hard tissue loss.

Box 1: A classification for managing maxillofacial wounds

This system focuses the surgeon's mind onto whether wounds can be closed primarily or whether the degree of hard and soft tissue loss means that later replacement and reconstruction is required. Any missing teeth should be accounted for as they may compromise the airway. Teeth lying in the airway adjacent to an endotracheal tube, buried in soft tissues or within the sinuses can be detected on CT scan.

The ready availability of CT scan with 3 dimensional reformatting improves clinical diagnosis. However, in Kandahar the patient is frequently passed from the CT scanner already intubated and into the operating theatre. It is important in the emergency setting to take time to methodically look through the CT scans before operating and record where shrapnel is and the precise sites and pattern of injury. In the finer bones of the midface, CT reconstructed images may fill in fractures that are undisplaced (Figures 1&2). Therefore, CT scans should not be relied upon alone and a thorough clinical evaluation of the common sites of fractures must be made prior to surgery. The assessment of the occlusion may be difficult in the orally intubated patient.

Initial Management

The initial management of facial injuries follows established military protocols. These protocols are similar to those of Advance Trauma Life Support [3] but differ in that the control of catastrophic, life threatening haemorrhage is dealt with first [4]. A cricthyroidotomy is the best way to establish a surgical airway in an emergency. A tracheostomy should only be done in theatres in a controlled environment and the flange sewn to skin to prevent dislodgement of the tube during evacuation. Maxillofacial haemorrhage is frequently stopped by local measures at the site of injury and temporary reduction and fixation of fractures. The face




Figure 1. Preoperative 3D CT image showing right zygomatic complex fracture (arrowed) orbital floor and nasal fractures.




Figure 2. Post-operative 3D CT image showing undisplaced Le fort 1 (fracture arrowed).

is rarely the site of torrential haemorrhage sufficient to be the sole contributor to hypovolaemic shock. However, when this does occur ligation of the ethmoidal vessels or the external carotid artery may be required. Patients with maxillofacial ballistic injuries should have a low threshold for placement of a tracheostomy tube. The likelihood of later airway swelling, further operations in a compromised patient and the need to correctly position the occlusion without the interference of an oral endotracheal tube may all necessitate a tracheostomy.

Once life saving measures have taken place, the patient requires resuscitation and initial surgical stabilisation [5]. The concept of damage control resuscitation and surgery applies to all severely injured oral and maxillofacial surgery patients and is vigorously applied at the Kandahar Role 3 Hospital. Damage control resuscitation aims to prevent the lethal triad of hypoxia, acidosis and coagulopathy by permissive hypotension and haemostatic resuscitation [6]. The goal of damage control surgery is to limit the physiological insult of surgery by carrying out the minimum amount of surgery in the shortest time to stabilize patients and prevent infection [7, 8]. Definitive surgery is done when the patient's condition has been optimized.

Oral and Maxillofacial Surgery

Early and aggressive debridement of wounds is required to prevent infection. This may involve the use of scrubbing brushes, pulsed lavage and copious irrigation. Surgical dermabrasion with a scalpel blade can be used to remove all debris that may cause subsequent wound tattooing which is difficult to correct. Deep penetrating wounds that are difficult to debride should be packed with iodine soaked gauze. To minimize unaesthetic scarring, superficial wounds that have been adequately cleaned are best closed primarily and rotation flaps may be used to close small defects. In high energy exchange ballistic trauma the wound is contaminated and the zone of tissue devitalisation around the missile tract will be unclear. These wounds should be cleaned and packed and serially debrided over several days until the vital wound margins have declared themselves [9]. Judgment is required to adequately clean the wound but not to strip vital periosteum or remove potentially viable tissue. In the early closure of soft tissue defects with rotational or advancement flaps thought should be given to possible later reconstruction if the flap breaks down. The rotational flap should not compromise the blood supply to a larger soft tissue flap. If it does it may be best to pack the area and close the defect once the tissues are seen to be clean and healthy.

Simple bony trauma can be repaired using conventional techniques of open reduction and internal fixation with craniofacial plates. However, caution must be exercised in using plates in contaminated wounds or wounds with associated large soft tissue defects. Primary reconstruction with bone plates and screws in the austere combat environment often yield poor results with subsequent soft tissue infections and plate exposures [10]. Inter maxillary fixation (IMF) using IMF bone screws [11] or arch-bars provide good reduction and often adequate immobilisation of mandibular fractures. Patients require wire cutters to be with them at all times to cut wire or elastics placed between the IMF screws or arch bars in case of vomiting which may compromise the airway. External fixators are robust and simple to apply. No foreign material is introduced at the fracture site and any soft tissue or bony defects may continue to be debrided with the fixator in place. External fixators are well suited to treated battlefield facial fractures and have been used frequently in Afghanistan. Combinations of plates, IMF, and external fixators can be used (Figures 3, 4 & 5).

Poor outcomes of ballistic facial injuries are often due to scarring and fibrous contracture in areas of severe tissue damage. Infection may have led to wound breakdown and increased the scarring and fibrosis. Regions of soft tissue avulsion closed over bony defects lack bone support and will contract, distorting the surrounding tissues. The resulting aesthetic and functional compromise may be severe. To optimise surgical outcomes, definitive surgery should be considered from the first operation. Infection must be prevented but wounds should be closed as early as possible to minimise scarring. Bones should be held in their correct position with IMF, external fixators or, where applicable, bone plates. Early bone repair should be done when all infection has been cleared and ideally within 3 to 4 weeks to minimize fibrosis and collapse of the soft tissue envelope [12].

NATO or western civilian personnel are evacuated back to their own Countries where reconstruction using free flaps,

osseointegrated implants and prosthesis, or osteogenic distraction techniques may be considered. In the austere environment of Afghanistan reconstructive surgery for local nationals should be kept as simple as possible in order to obtain the best results. This may involve scar revision, rotational flaps and bone grafts.

Rehabilitation with adequate nutrition, good oral hygiene and physiotherapy is an important part of treatment. Tongue spatulas may be used between the teeth to stretch facial scars and prevent trismus (Figures 6 & 7).



Figure 3. Preoperative 3D CT image showing severe ballistic complex comminuted facial fractures.



Figure 4. Treatment of mid-face fractures with an external fixator, intermaxillary fixation and mandibular palates.




Figure 5. Preoperative 3D CT image showing severe ballistic complex comminuted facial fractures.



Figure 6. Facial scarring leading to trismus.



Figure 7. Treatment with tongue spatulas to stretch scars and improve mouth opening.

Body Armour and Measures for Facial Injury Prevention

The use of body armour has reduced the overall number of deaths and torso injuries but has increased the relative proportion of wounds to the extremities including the head and neck [2]. Oral and maxillofacial surgeons often saw patients with damaged protective glasses that had prevented penetrating eye injuries from shrapnel. Further work is needed on facial armour that can protect the face without limiting visual fields and mobility. In vehicles, soldiers on 'top cover' frequently present with blast injuries to their exposed faces. This particular group would gain maximum benefit from facial armour. Military personnel within troop carriers rarely wear seat belts as they may not fit over body armour and are thought to restrict rapid evacuation. Hence, when the vehicle is hit by an IED the soldiers are thrown around inside the vehicle and incur limb and facial lacerations and fractures. Quick release seat belts that are able to be fastened over bulky body armour would help prevent these injuries and this is an area of ongoing research.

Conclusions

The role of military oral and maxillofacial surgeons within the Role 3 Hospital at Kandahar is established. With the high workload, a wealth of experience in the management of ballistic trauma can be gained by surgeons in just a few months. Since returning from deployment the British Oral and Maxillofacial Surgery Cadre has held regular clinical meetings. From the knowledge acquired, several teaching initiatives have taken place. These include presentations on the Military Operational Surgical Training (MOST) course at the Royal College of Surgeons of England and for the Norman Rowe International Education Foundation work in post conflict countries. The MOST course is held over 5 days twice a year for military medical staff and aims to transfer skills between surgical specialities and give an update on current surgical techniques used in the management of battlefield trauma.

References

1. Rustemeyer J, Kranz V, Bremerich A. Injuries in combat from 1982-2005 with particular reference to those to the head and neck: A review. *Br J Oral Maxillofac Surg* 2007; **45**: 556-560
2. Ramasamy A, Harrison SE, Stewart MPM, Midwinter M. Penetrating missile injuries during the Iraqi insurgency. *Ann R Coll Surg Engl* 2009; **91**: 551-558.
3. American College of Surgeons committee on Trauma. *Advanced Life Support for doctors*. 8th edn.
4. Hodgetts TJ, Mahoney PF, Russell MQ, Byres M. ABC to < C > ABC: redefining the military trauma paradigm. *Emerg Med J* 2006; **23**: 745-746.
5. Gibbons AJ, Patton DW. Ballistic Injuries of the Face and Mouth in war and Civil Conflict. *Dental Update* 2003; **30**: 272-278
6. Jansen JO, Thomas R, Loudon MA, Brooks A. Damage control resuscitation of patients with major trauma. *Br Med J* 2009; **338**: 1436-1440
7. Blackburne LH. Combat damage control surgery. *Crit Care Med* 2008; **36**: S304-S310.
8. Bowley DM, Barker P, Boffard KD. Damage control surgery- concepts and practice. *J R Army Med Corps* 2000; 176-182
9. Robertson BC, Manson PN. High-energy ballistic and avulsive injuries. A management protocol for the next millennium. *Surg Clin North Am*. 1999; **79**:1489-1502.
10. Gokel T. Improvised Explosive Devices and the Oral and Maxillofacial Surgeon. *Oral Maxillofacial Surg Clin N Am* 2005; **17**: 281-287
11. Gibbons AJ, Baden JM, Monaghan AM, Dhariwal DK, Hodder SC. A drill-free bone screw for intermaxillary fixation in military casualties. *J R Army Med Corps* 2003; **149**: 30-32
12. Epeley BL, Coleman JJ. Reconstruction of Large Hard and Soft Tissue Defects of the Face. In: *Maxillofacial trauma and Esthetic Reconstruction*, Ward Booth P, Epeley B, Schmeltzseisen R (Eds). Churchill Livingstone, Elsevier; 2003